

The surface density profile of NGC 6388: a good candidate for harboring an intermediate-mass black hole ¹

B. Lanzoni¹, E. Dalessandro^{1,2}, F.R. Ferraro¹, P. Miocchi³, E. Valenti⁴, R.T. Rood⁵

¹ *Dipartimento di Astronomia, Università degli Studi di Bologna, via Ranzani 1, I-40127 Bologna, Italy*

² *ASI, Centro di Geodesia Spaziale, contrada Terlecchia, I-75100, Matera, Italy*

³ *Dipartimento di Fisica, Università di Roma “La Sapienza,” P.le A. Moro 5, I-00185 Roma, Italy*

⁴ *European Southern Observatory, Alonso de Cordova 3107, Vitacura, Santiago, Chile*

⁵ *Astronomy Department, University of Virginia, P.O. Box 400325, Charlottesville, VA, 22904*

24 July, 07

ABSTRACT

We have used a combination of high resolution (*HST* ACS-HRC, ACS-WFC, and WFPC2) and wide-field (*ESO*-WFI) observations of the galactic globular cluster NGC 6388 to derive its center of gravity, projected density profile, and central surface brightness profile. While the overall projected profiles are well fit by a King model with intermediate concentration ($c = 1.8$) and sizable core radius ($r_c = 7''.2$), a significant power law (with slope $\alpha = -0.2$) deviation from a flat core behavior has been detected within the inner $1''$. These properties suggest the presence of a central intermediate mass black hole. The observed profiles are well reproduced by a multi-mass isotropic, spherical model including a black hole with a mass of $\sim 5.7 \times 10^3 M_\odot$.

Subject headings: Globular clusters: individual (NGC6388); stars: evolution – black hole physics

¹Based on observations with the NASA/ESA *HST*, obtained at the Space Telescope Science Institute, which is operated by AURA, Inc., under NASA contract NAS5-26555. Also based on WFI observations collected at the European Southern Observatory, La Silla, Chile.

1. INTRODUCTION

The surface brightness (SB) and the projected density profiles of the vast majority of globular clusters (GCs) are well reproduced by a family of simple models characterized by an extended isothermal core and a tidally truncated envelope—the so-called King models (King 1966). However a fraction ($\sim 15\text{--}20\%$, see Djorgovski & King (1986)) of galactic GCs deviate significantly from this behavior. The projected density profiles of these clusters do not exhibit an extended core, showing instead a power law behavior $\Sigma(r) \propto r^\alpha$ with α ranging from -0.8 to -1.0 . This feature has been thought to arise from the dynamical evolution of stellar systems that have experienced the collapse of the core. These are called post-core collapse clusters, hereafter PCC).

However, other processes can affect the shape of a star cluster density profile; among these the existence of an Intermediate Mass Black Hole (IMBH) in the central region has recently received attention. Interestingly enough, detailed collisional N -body simulations (Baumgardt et al., 2005, hereafter BMH05; Trenti et al. 2007) and theoretical arguments (Heggie et al. 2007) have shown that the presence of a IMBH yields quite a different SB profile than core collapse does. These studies indicate that in an initially dense cluster a IMBH gives rise to a strong expansion of the central region that, in turn, leads to a quasi-steady configuration resembling that of a medium concentration cluster with a core-like profile. Thus, the clusters most likely to harbor IMBHs are those having the appearance of normal King model clusters except in the very central regions where a power law deviation from a flat behavior is expected. The exponent of this power law is predicted to be significantly lower ($\alpha \sim -0.2$) than in the PCC case (BMH05, Miocchi 2007). Small departures from a King model have been observed in the projected density profile of a few GCs (included NGC6388) by Noyola & Gebhardt (2006, hereafter NG06).

The confirmation of the existence of IMBHs and an estimate of their frequency would be important for a number of astrophysical problems like the formation processes of super massive BH in galaxies, super-Eddington X-ray sources in extragalactic globular clusters (Sivakoff et al. 2007), the origin of ultraluminous X-ray sources (Miller 2003; Fabbiano 2006). Despite of their potential importance, the existence of IMBH in GCs is still an open question. For instance, the evidence for an IMBH in M15 reported by van der Marel et al (2002) and Gerssen et al (2002) has been questioned by Baumgardt et al (2003a). Baumgardt et al (2003b) also question evidence for an IMBH in G1 in M31 (Gebhardt, Rich & Ho 2002) (but see Gebhardt, Rich & Ho (2005) and the recent findings by Ulvestad et al (2007) and Green & Ho, 2007).

Here we present accurate surface density and SB profiles obtained with a combination of high-resolution and wide-field observations of the galactic globular cluster NGC 6388, which

a number of authors (BMH05; NG06; Drukier & Bailyn 2003, hereafter DB03; Miocchi 2007) have indicated as a prime candidate to harbor an IMBH. These profiles nicely match the theoretical models of a cluster with a $5.7 \times 10^3 M_\odot$ BH: having an extended core, intermediate concentration, but also significant deviations from a flat core distribution in the innermost cluster regions ($r \lesssim 1''$).

2. THE DATA

In this paper we make use of a combination of high-resolution and wide-field photometric data sets, obtained with WFPC2 and ACS on board *HST*, and with WFI at ESO-2.2 m, respectively. A detailed description of the observations, photometric reduction and astrometry procedures of the data obtained with WFPC2, ACS-WFC, and WFI is given in a companion paper discussing the Blue Straggler Star and Horizontal Branch populations (Dalesandro et al. 2007). Here we use only the optical (B , V , I) samples from the entire multi-wavelength data set. These have been all homogenized and transformed to the Johnson magnitude system. All the star positions have been placed on the absolute astrometric system using several hundred astrometric reference stars from the new astrometric 2-MASS catalogue², following the procedure described, e.g., in Lanzoni et al. (2007), with a final astrometric accuracy of the order of $\sim 0''.3$ both in RA and Dec.

Additional images obtained with the ACS-HRC have been retrieved from the ESO/ST-ECF Science Archive. These data sample the cluster central region with a field of view (FoV) of $26'' \times 29''$ and a spatial resolution of $0''.027 \text{ pix}^{-1}$. The HRC data were obtained through filters $F555W$ (V) and $F814W$ (I), with total exposure times of 620 and 3070 s, respectively. After corrections for geometric distortions and effective flux (Sirianni et al. 2005), the photometric analysis was performed by using SExtractor (Bertin & Arnouts 1996), adopting a fixed aperture radius of 4 pixels ($0''.108$). The sample has then been astrometrized and photometrically calibrated by using the stars in common with the ACS-WFC catalog. The color-magnitude diagrams based on the data from all four data sets are shown in Fig. 1.

3. CENTER OF GRAVITY

Given the absolute positions of individual stars in each catalog, the center of gravity, C_{grav} , of NGC 6388 has been determined by averaging the coordinates α and δ of all stars

²Available at <http://irsa.ipac.caltech.edu>.

detected in the highest resolution data set (the HRC sample). In order to correct for spurious effects due to incompleteness in the very inner regions of the cluster, we considered only stars brighter than $V = 20$ (roughly corresponding to the sub-giant branch of the cluster). By following the iterative procedure described in Montegriffo et al. (1995, see also Ferraro et al. 2004), the center of gravity is located at $\alpha_{J2000} = 17^{\text{h}} 36^{\text{m}} 17^{\text{s}}.23$, $\delta_{J2000} = -44^{\circ} 44' 7''.1$, with an uncertainty of $0''.3$ in both α and δ . A careful examination of field inside the core radius shows that our determination of the center is biased neither by the presence of very bright stars, nor of a star clump. The derived C_{grav} is located $\sim 2''.6$ southeast ($\Delta\alpha = 3''.4$, $\Delta\delta = -1''.1$) of that derived by Djorgovski & Meylan (1993) using the surface brightness distribution. An accurate comparison with the center adopted by NG06 is not possible since the value listed in their Table 1 is just referred to the world coordinate system of a specific WFPC2 image, however a visual inspection suggests that their center is located $\sim 0''.5$ SE of ours.

4. PROJECTED DENSITY AND SURFACE BRIGHTNESS PROFILES

We have determined the projected density profile of NGC 6388 using direct star counts over the entire cluster radial extent, from C_{grav} out to $\sim 1400'' \sim 23'$. This distance is significantly larger than the expected cluster extension ($r_t = 372''$, Harris 1996). In order to avoid spurious effects due to possible incompleteness, only stars brighter than $V = 20$ have been considered (see Dalessandro et al 2007). There are more than 58,000 stars in the entire (i.e., the combination of ACS, WFPC2 and WFI) photometric data set. Following the procedure described in Ferraro et al. (1999a), we have divided the sample in 40 concentric annuli, each centered on C_{grav} . Each annulus has been split into an adequate number of sub-sectors. The number of stars lying within each sub-sector was counted, and the star density was obtained by dividing these values by the corresponding sub-sector areas. The stellar density in each annulus was then obtained as the average of the sub-sector densities, and the uncertainty in the average values for each annulus was estimated from the variance among the sub-sectors. The radius associated with each annulus is the mid-point of the radial bin. The outermost ($r \gtrsim 7'$) measures have an almost constant value and their average has been used to estimate the Galaxy contamination to be ~ 56 stars arcmin $^{-2}$. Subtracting this background yields the the profiles shown in Figs. 2.

If the innermost ($r < 1''$) points are excluded, the density profile is well fit all over the entire extension by an isotropic, single-mass King model with a core of $r_c = 7''.2$ and an intermediate concentration ($c = 1.8$). These values are similar to those quoted by Trager et al. (1995; $r_c = 7''.4$ and $c = 1.7$), Harris (1996; $r_c = 7''.2$ and $c = 1.7$), and McLaughling &

van der Marel (2005; $r_c = 7''.8$ and $c = 1.71$).³ In the inner $\sim 1''$ the observed profile shows an indication of a deviation from a flat core behavior. With only 7 stars in the innermost bin, the statistical error of the inner bin ($0'' \leq r < 0''.3$) is relatively large. With star counts alone the exact amount of the deviation from the flat core cannot be reliably estimated.

Exploiting the exceptional high resolution of ACS-HRC images we have computed the SB profile by direct aperture photometry to more accurately determine the inner shape of the cluster profile. In the innermost region ($r < 1''$) we have used two sets of annuli stepped by $0''.3$ and $0''.5$, respectively. The SB values have been computed as the sum of the counts in each pixel, divided by the number of pixels in any given annulus. The counts have then been converted to a magnitude scale and calibrated by using a relation that links the “instrumental” magnitude to the calibrated one (obtained by performing aperture photometry for a number of high S/N isolated stars). The resulting SB profile for the innermost $10''$ from the center is shown in Figure 3. A steepening of the profile at $r \lesssim 1''$ is clearly apparent, in agreement with what we found above for the surface density distribution. A linear fit to the inner points suggest that the slope of the profile is 0.6 ± 0.06 in the $\log \Sigma - \log r$ plane. In terms of the surface luminosity density $I(r)$, if $I \propto r^\alpha$ we find $\alpha \simeq -0.23$. This is steeper (but still marginally consistent within the errors) than the slope $\alpha = -0.13 \pm 0.07$ derived from the analysis of WFPC2 images by NG06. The N06 profile is shown for comparison in Figure 3, as can be seen their profile is fully compatible with our data in the common region. The use of high resolution images allow us to probe the innermost region of the cluster where most of the deviation from a flat behavior occurs⁴.

5. DISCUSSION

The properties of the projected density and SB profiles derived in the previous section for NGC 6388 are not those of a cluster which has experienced the core collapse. Instead they are just what BMH05 suggest as the signatures of a cluster harboring an IMBH in its center: (i) a typical King profile with intermediate concentration ($c = 1.8$) in the external regions, (ii) a sizable core, and (iii) a inner logarithmic slope $\alpha \sim -0.2$. These features have been recently confirmed by the predictions of a self-consistent parametric model that includes the

³Note that the higher concentration quoted in the present paper implies a ($\sim 20\%$) larger tidal radius ($r_t = 454''$) than previously determined.

⁴Even the small difference in the center determination can play a role. Simulations have shown that even an offset of only a few $0''.1$ is sufficient to flatten the profile. An additional difference in the slope determination might arise from the different approach used by NG06, who removed the bright stars and did not fit a power law to the data but instead calculated the derivative of the smooth central profile.

presence of a central IMBH (Miocchi 2007). This model consists of a multi-mass isotropic, spherical King model, which has been extended inside the region of gravitational influence of the BH via the inclusion of the Bahcall & Wolf (1976) phase-space distribution function.

In order to further support the case for an IMBH in the center of NGC 6388, we have used this model to reproduce the observed density and SB profiles. A Salpeter mass function ($dN \propto m^{-1.35} d\log m$) is assumed in the model and seven discrete mass bins are used to approximate the continuum mass spectrum of the real cluster. The stellar mass range in the interval from 0.3 to $0.9 M_\odot$ (here and thereafter, where not specified, masses are measured in $M_\odot$) equally subdivided in 6 bins 0.1 wide. These were populated with main sequence stars. In addition the bins $[0.5, 0.6]$ and $[0.7, 0.8]$ include WD populations with mass 0.55 and 0.75 , coming from progenitors with mass, respectively, in $[0.9, 1.5]$ and $[1.5, 4]$ ranges. The seventh mass bin contains a massive WD population with $m = 1.2$, hypothesized as remnants of stars with mass $4-8$. The $[0.8, 0.9]$ bin has $\langle m \rangle = 0.84$ and contains the TO stars, plus giants and HB stars. The light-to-mass ratios were taken to be $\{4.9 \times 10^{-3}, 10^{-2}, 2.3 \times 10^{-2}, 6.5 \times 10^{-2}, 0.19, 10, 0\}$ corresponding to the bins ordered in increasing average mass. The velocity dispersion of the seven components is constrained by the requirement of complete energy equipartition at the border of the BH influence region (see Miocchi 2006; Miocchi 2007, for details), where the adimensionalised potential W_{BH} , along with the ratio between the BH mass (M_{BH}) and the cluster mass (M), determine the form of the generated profiles. Besides of the various scale parameters, W_{BH} and M_{BH} are adjusted to obtain the best fit to the observed profiles. To do this, we conservatively include only data from the central $100''$ in order to avoid possible spurious effects which might affect the most external points of the SB profiles because of the field contamination subtraction. The best fit to the observed SB profile is then found for $r_c = 7''.2$, $c = 1.8$, $W_{\text{BH}} = 11.5$ and $M_{\text{BH}} = 2.2 \times 10^{-3} M$ (yielding $P(\chi^2 > \chi_{\text{fit}}^2) > 99\%$). The level of confidence remains above 97% for an IMBH mass in the range $2.1-2.4 \times 10^{-3} M$. The r_c and c values are consistent with the value deduced by the parametric fit of the single-mass King model mentioned above. The results of this parametric fit to the projected density and the SB profiles are shown in Figs. 2 and 3 respectively.

By assuming the total cluster luminosity $V_t = 6.72$ (Harris 1996), and the distance modulus $(m - M)_V = 16.59$ (Dalessandro et al. 2007)⁵, we estimate a total mass of $2.6 \times 10^6 M_\odot$ for NGC 6388, yielding $M_{\text{BH}} \simeq 5.7 \times 10^3 M_\odot$.⁶

⁵The distance has been obtained differentially with respect to 47 Tucanae, by assuming the distance scale by Ferraro et al. (1999b).

⁶Note that this mass is well within the range of values ($2.5-10 \times 10^3$) derived from the $M_{\text{BH}} - \sigma$ relation (Ferrarese & Merritt 2000; Gebhardt et al. 2000; see for example equation (10) of DB03), by assuming the

While the central IMBH is a possible explanation of the shape of the observed profiles, one might question whether this result is unique. In fact, a central concentration of massive remnants (like white dwarfs and neutron stars) has been proposed as an alternative explanation in the case of M15 (van den Bosch et al 2006). However, we have found that a multi-mass King model including a population of such remnants but without central BH is unable to reproduce the observed slope of the profiles in the core region. Since our evidence is based exclusively from the shape of the density profile, the presence of a IMBH at the center of NGC 6388 is still debatable. Accurate kinematical studies of the motion of stars in the central region of the cluster are needed to solidify the case.

The region in which to seek for the possible kinematical signatures of a BH is very small. The self-consistent model here employed generates a projected velocity dispersion profile that shows a sharp rise from the “isothermal plateau” to a purely Keplerian behavior at $r \sim 0''.16$ (i.e., $\sim 0.02 r_c$). A more promising path to detect the kinematic signature of a BH is by proper motion measurements (DB03). Some stars should move with anomalously high velocities under the influence of the BH. In order to estimate the number of these stars, we first need to evaluate the BH radius of influence r_h . A crude estimate of r_h is given by: $r_h = G M_{\text{BH}} / \sigma_0^2$, where σ_0 is the velocity dispersion outside r_h . By assuming $\sigma_0 = 18.9 \text{ km s}^{-1}$ (Pryor & Meylan 1993), we find $r_h \simeq 0.07 \text{ pc}$, corresponding to $1''.1$ with the cluster distance ($d = 13.18 \text{ kpc}$; Dalessandro et al. 2007). By using equation (7) of DB03, it is possible to estimate the number of stars (N), measurable through proper motions studies, having velocities 3 times the cluster velocity dispersion outside r_h : $N = 0.27 \Sigma_0 r_h^2$, where $\Sigma_0 r_h^2$ is the number of stars within r_h . This relation suggests that $\sim 27\%$ of stars within r_h are expected to show anomalously high velocity. We can directly derive this number from the HRC images. Adopting the value of the cluster center presented above we count 28 relatively bright ($V < 19$) stars within $r_h = 1''.1$ (out of a total of ~ 85 stars detected down to $V \sim 22$), corresponding to a total of ~ 7 high-velocity stars. This estimate shows that the presence of $\sim 5 \times 10^3 M_\odot$ BH in the center of NGC 6388 can in principle be kinematically confirmed in the near future through accurate proper motion studies or radial velocity measurements with *Adaptive Optic* supported instruments. However, these measurements can be quite challenging. According to Figure 1 of DB03, the high velocity stars are expected to be mainly confined within $0.4 r_h$ (only $\sim 0''.4$ from the center). They would have speeds of order 60 km s^{-1} which, given projection effects, would give tangential velocities of order 20 km s^{-1} . The distance derived by Dalessandro et al. (2007) and adopted here is 25% larger than that from Harris (1996), so the resulting proper motions would be $\sim 0.3 \text{ mas yr}^{-1}$. Given the current estimate for the proper motion measurement error of roughly 0.3 mas , a baseline of

low ($\beta = 4.02$) and the high ($\beta = 4.65$) exponents, respectively.

at least 3–5 yr would be required.

This research was supported by contract ASI-INAF I/023/05/0, PRIN-INAF2006 and by the Ministero dell’Istruzione, dell’Università e della Ricerca. ED is supported by ASI. RTR is partially supported by STScI grant GO-10524. This research has made use of the ESO/ST-ECF Science Archive facility which is a joint collaboration of the European Southern Observatory and the Space Telescope - European Coordinating Facility.

REFERENCES

- Bahcall J.N., Wolf R.A, 1976, ApJ, 209, 214
- Baumgardt H., Makino J., Hut P., 2005, ApJ, 620, 238 (BMH05)
- Baumgardt, H., Hut, P., Makino, J., McMillan, S., Portegies Zwart, S., 2003a, ApJ, 582, L21
- Baumgardt, H., Makino, J., Hut, P., McMillan, S., Portegies Zwart, S., 2003b, ApJ, 589, L25
- Bertin, E., & Arnouts, S. 1996, A&AS, 117, 393
- Djorgovski, S., & Meylan, G. 1993, Structure and Dynamics of Globular Clusters. Proceedings of a Workshop held in Berkeley, California, July 15-17, 1992, to Honor the 65th Birthday of Ivan King. Editors, S.G. Djorgovski and G. Meylan; Publisher, Astronomical Society of the Pacific, Vol. 50, 325
- Djorgovski, S., & King, I.R., 1986, ApJ, 305, L61
- Dalessandro E., et al., 2007 in preparation
- Drukier, G.,A., Bailyn, C.D., 2003, ApJ, 597, L125 (DB03)
- Fabbiano, G., 2006, ARA&A, 44, 323
- Ferrarese, L., Merritt, D., 2000, ApJ, 539, L9
- Ferraro, F. R., Paltrinieri, B., Rood, R. T., & Dorman, B. 1999a, ApJ 522, 983
- Ferraro F. R., Messineo M., Fusi Pecci F., De Palo M. A., Straniero O., Chieffi A.,& Limongi M. 1999b, AJ, 118, 1738

- Ferraro, F. R., Beccari, G., Rood, R. T., Bellazzini, M., Sills, A., & Sabbi, E. 2004, *ApJ*, 603, 127
- Gebhardt, K., et al., 2000, *ApJ*, 539, L13
- Gebhardt, K., Rich, R.M., Ho, L.C. , 2002, *ApJ*, 578, L41
- Gebhardt, K., Rich, R.M., Ho, L.C. , 2005, *ApJ*, 634, 1093
- Gerssen, J., van der Marel, R.P., Gebhardt, K., Guhathakurta, P., Peterson, R. C.; Pryor, C., 2002, *ApJ*, 124, 3270
- Greene, J.E., Ho, L.C., 2007, *ApJ*, 656, 84
- Harris, W.E. 1996, *AJ*, 112, 1487
- Heggie D. C., Hut P., Mineshige S., Makino J., Baumgardt H., *PASJ Letters*, *PASJ Vol. 59* (astro-ph/0611950)
- King I.R., 1966, *AJ*, 71, 64
- Lanzoni, B., Dalessandro, E., Ferraro, F. R., Mancini, C., Beccari, G., Rood, R. T., Mapelli, M., & Sigurdsson, S. 2007, *ApJ*, 663, 1040
- McLaughlin, D. E., & van der Marel, R. P. 2005, *ApJS*, 161, 304
- Miller, C. 2003, Dynamics and Evolution of Dense Stellar Systems, 25th meeting of the IAU, Joint Discussion 11, 18 July 2003, Sydney, Australia, 11,
- Miocchi P., 2006, *MNRAS*, 366, 227
- Miocchi, P. 2007, in press of *MNRAS*(astro-ph/0702479)
- Montegriffo, P., Ferraro, F. R., Fusi Pecci, F., & Origlia, L. 1995, *MNRAS*, 276, 739
- Noyola, E., Gebhardt, K., 2006, *AJ*, 132, 447 (NG06)
- Pryor C., & Meylan G., 1993, Structure and Dynamics of Globular Clusters. Proceedings of a Workshop held in Berkeley, California, July 15-17, 1992, to Honor the 65th Birthday of Ivan King. Editors, S.G. Djorgovski and G. Meylan; Publisher, Astronomical Society of the Pacific, Vol. 50, 357
- Sivakoff, G. R., et al. 2007, *ApJ*, 660, 1246
- Sirianni, M., et al. 2005, *PASP*, 117, 1049

- Trager, S.C., Djorgovski, S., & King I.R., 1993, Structure and Dynamics of Globular Clusters. Proceedings of a Workshop held in Berkeley, California, July 15-17, 1992, to Honor the 65th Birthday of Ivan King. Editors, S.G. Djorgovski and G. Meylan; Publisher, Astronomical Society of the Pacific, Vol. 50, 347
- Trager, S.C., King I.R. & Djorgovski, S., 1995, AJ, 109, 218
- Trenti M., Ardi E., Mineshige S., Hut P., 2007, MNRAS, 374, 857
- Ulvestad, J.S., Greene, J.E., Ho, L.C., 2007, ApJ, 661, L151
- van den Bosch, R., de Zeeuw, T., Gebhardt, K., Noyola, E., van de Ven, Glenn, 2006, ApJ, 641, 852
- van der Marel, R.P., Gerssen, J., Guhathakurta, P., Peterson, R.C., Gebhardt, K., 2002, AJ, 124, 3255




Fig. 1.— Color Magnitude diagrams for the four data sets used. The WFI data are dominated by field contamination.




Fig. 2.— Observed surface density profile obtained by star counts from the combined photometric data-set: ACS-HRC (*filled circles*), WFPC2 (*triangles*), ACS-WFC (*squares*), and WFI (*empty circles*). The background level (see Sect. 4) has been subtracted. The single-mass King model that best fits the profile excluding the innermost ($r \lesssim 1''$) points is shown as a dashed line. It is characterized by a sizable core radius ($r_c = 7''.2$), and an intermediate concentration ($c = 1.8$). The solid line shows the profile of the model (see Sect. 5) including a $5.7 \times 10^3 M_\odot$ BH in the cluster center. The profile from Trager et al (1995) is shown (*dotted line* for comparison).




Fig. 3.— Surface brightness profile derived from the ACS-HRC images within $10''$ from the cluster center. *Dots* refer to a radial binning of $0''.5$, triangles sample the inner $1''$ by steps of $0''.3$. The *solid line* shows the profile of the model obtained by including a $5.7 \times 10^3 M_\odot$ BH (the same as in Fig. 2). The *short-dashed line* corresponds to the King model shown in Fig. 2. $\beta = 0.6$ is the slope of the linear best fit (see the *long-dashed line*) to the innermost points. The profile obtained by NG06 is also shown (see the *dotted line* for comparison).